issued by the Jacksonville office and promptly distributed by the local offices in cooperation with all communication

Disturbance of October 18-21.—The first indications of a definite circulation in connection with this slight tropical disturbance were noted about 100 miles off the west Florida coast on the night of October 18-19. It had developed from a wave of low pressure and squally weather that had moved across the southern Bahamas and through the Florida Straits a few days previously, and for which small-craft warnings had been issued on the morning of October 17.

On the morning of October 19, a ship, 50 miles west of Tampa, reported a west-southwest wind of 45 miles per hour and a barometer reading of 1,002.4 millibars (29.60 inches). From this location the storm center then moved

north-northwestward and passed inland at Cedar Key, about noon of the 20th, where the lowest barometer reading was 1,005.8 millibars (29.70 inches).

Available observations show no winds over moderate gale force (40 to 50 miles per hour) during the progress of this storm. Torrential rains (10 to 15 inches) occurred at several points near the center as the disturbance stalled and dissipated over northern Florida.

A report from Ocala lists the death of a 6-weeks-old infant, hurled 100 feet from its basket, and injury to both parents when their house was demolished by high winds. This report indicates the formation of a small and short-lived tornado rather than any highly destructive winds resulting directly from the tropical disturbance. All other reported damage resulted from flooding due to the excessive rains.

METEOROLOGICAL AND CLIMATOLOGICAL DATA FOR OCTOBER 1941

[Climate and Crop Weather Division, J. B. KINCER in charge]

AEROLOGICAL OBSERVATIONS

By Homer D. Dyck

Departures from normal surface temperature for October followed nearly the same pattern as in the previous month, temperatures being above normal generally over the eastern half of the country with the exception of an area in the extreme Northeast, and below normal over the western half with the exception of the far Northwest and a narrow strip along the Pacific Coast. Plus departures ranged from 6° to 8° F. over much of the South, while minus departures were small.

At 1.500 meters above sea level the 5 a.m. resultant winds for October were from directions to north of normal over the Lake region and the extreme Northeast and over much of the Plateau region and the far Northwest, while they were from south of normal at this level over much of the rest of the country. At 3,000 meters the morning resultant winds for October were from directions south of normal over most of the country with the exception of three stations near the Pacific coast and one near the Lake region, which had resultant winds to north of normal. At 5,000 meters a good comparison of the 5 p. m. resultant winds with the corresponding 5 a.m. normals was not possible over the northern Great Plains and the Lake region. Elsewhere in the country the October afternoon resultants at this level were from directions to south of the corresponding morning normals generally, except over the Pacific Coast States and over a few stations in the East.

At both the 1,500- and 3,000-meter levels resultant velocities were below normal over the northern Plateau region, the central Mississippi Valley and an area in the southeastern States. At 5,000 meters the 5 p. m. resultant velocities were decidedly higher than corresponding morning normals everywhere except in the Northwest.

Corresponding to the similarity of the October surface temperature departure pattern to that of September, there is also a marked similarity in the departures from normal resultant wind directions for the two months. The same general characteristics are evident in both months, i. e., the turning to southward of normal generally over the eastern half of the country, and the opposite shift from normal over the western half. During October, however, the area in the west where turning to northward of normal took place, was not nearly as extensive as in September thus accounting perhaps for the smaller minus temperature anomalies over that area during October.

When the 5 p. m. resultant directions are compared to the corresponding 5 a. m. resultant directions, no well marked areas of similar wind shift are evident. It may be noted, however, that a shift to southward during the day occurred at considerably more stations than a shift to northward.

At 1,500 meters the 5 p. m. resultant velocities were smaller than the corresponding 5 a. m. velocities over the extreme northern States and over the Middle Atlantic States and the South, while the afternoon velocities were larger than the morning velocities elsewhere generally. At 3,000 meters the p. m. resultant velocities were higher than corresponding a. m. velocities everywhere except over a small area in the southern Plateau region and California where they were slightly below the morning velocities.

The upper-air data discussed above are based on 5 a.m. (E. S. T.) pilot balloon observations (charts VIII and IX) as well as on observations made at 5 p.m. (table 2 and charts X and XI).

Radiosonde and airplane stations located in the southern part of the country recorded on the average the highest mean monthly pressures at each of the several standard levels from 2,000 to 18,000 meters. The highest mean monthly pressure occurred over both Atlanta and Charleston at the 2,000- and 2,500-meter levels, over Atlanta at 3,000 and 4,000 meters, and over Atlanta, Brownsville, and Miami at 5,000 meters. At 6,000 meters Atlanta, Brownsville, Miami, and San Antonio recorded the highest mean monthly pressure while the highest occurred over Brownsville, Lake Charles, Miami, and San Antonio at 7,000 meters, over Brownsville and Miami at 8,000 meters, over Miami and San Antonio at 9,000 meters, and over Brownsville, Miami, and San Antonio at 10,000 meters. At 11,000 meters the highest mean monthly pressure occurred over Miami and San Antonio, while at 12,000 meters pressures averaged highest over Brownsville, Miami, and San Antonio. San Antonio recorded the highest mean monthly pressure from 13,000 to 18,000 meters inclusive. Atlanta also recorded the same maximum, however at 16,000 meters. The lowest mean monthly pressure occurred over both Portland, Maine. and Sault Ste. Marie from 2,000 to 6,000 meters inclusive, while at 7,000 and 8,000 meters the lowest occurred over Portland, Maine, Sault Ste. Marie, Seattle, and Spokane, and at 9,000 meters it occurred at Spokane and Seattle. Spokane recorded the lowest mean monthly pressure at

levels from 10,000 to 16,000 meters inclusive, with Seattle also recording the minimum at the 14,000, 15,000, and 16,000 meter levels. Portland, Maine, also recorded the minimum at 14,000 and 16,000 meters. Pressures averaged lowest over Spokane at 17,000 and 18,000 meters. Mean monthly pressures at Alaskan stations were lower than mean minimum pressures over the United States at corresponding levels from 2,000 to 16,000 meters, inclusive.

Mean surface pressures for October were higher than those for September over most of the United States. At standard levels above the surface, however, only the Northwest recorded higher pressures than last month and these only up to about 7,000 meters, while the rest of the country recorded substantial decreases from last month at all standard levels. These decreases amounted to as much as 9 mb. over Portland, Maine, from about 5,000 to 10,000 meters. All Alaskan stations reported substantial decreases in pressure at all levels. Pressure gradients this month were less steep than last month over the Northwest, but were steeper in general elsewhere over the country. The steepest upper level pressure gradient for October occurred between Sault Ste. Marie and Detroit at the 8,000 meter level where there was a change of 1mb. pressure for each 38 miles of horizontal distance between the two stations.

The mean temperatures for October were considerably lower than September's at most levels up to and including 13,000 meters. Above this level scattered exceptions occurred over the southern Plateau region and the Lake

region.

Mean temperatures for October 1941 were considerably higher than those for October 1940 over most of the eastern half of the country up to and including 11,000 meters and somewhat below October 1940 over the western half up to about 7,000 meters. From 7,000 to 19,000 meters, temperatures over the western half were generally above last year's and from 13,000 to 19,000 meters temperatures over the eastern half of the United States were generally lower than last year's.

At 1,000 meters mean temperatures for October were above normal east of the Rocky Mountains with the exception of the central Great Plains and the extreme Northeast where they were slightly below normal. Elsewhere they were below normal at this level. At the 3,000- and 5,000-meter levels mean temperatures were below normal west of the Rocky Mountains and in the extreme Northeast and considerably above normal eslewhere.

The mean relative humidities for the month at the 1,000-, 3,000-, and 5,000-meter levels were considerably above normal over most stations. Marked plus departures occurred over San Antonio, Oklahoma City, and St. Louis at the 1,000- and 3,000-meter levels.

The altitude at which the mean monthly temperature of 0° C. for October occurred varied from the lowest (2,500 meters) over Seattle, Sault Ste. Marie, and Portland, Maine, to the highest (4,900 meters) over Brownsville, Tex. The level at which, on the average, freezing conditions occurred was lower this month than last over all of the United States with the exception of Pensacola, where it was slightly higher. This level was much lower than it was last month over the extreme Northeast, being 1,600 meters lower over Portland, Maine.

The lowest free-air temperature recorded during the month over the United States was -86.3° C. $(-123.3^{\circ}$ F.). This temperature occurred over Miami, Fla., on the morning of October 6, at an altitude of 16,300 meters (about 10.1 miles) above sea level. The lowest temperature for the month over San Juan was -82.0° C. $(-115.6^{\circ}$ F.) observed at 17,300 meters (about 10.8 miles) above sea

level on October 12.

Table 3 shows the maximum free-air wind velocities for various sections of the United States during October as determined by pilot balloon observations. The highest observed wind velocity for the month was 77 m. p. s. (172 miles per hour) observed over Great Falls, Mont., on October 2. This wind was blowing from the north at an elevation of 10,220 meters (about 6.4 miles) above sea level.

The highest October wind velocity observed during the last 5 years in the free-air layer from the surface to 2,500 meters was 46.8 m. p. s. (105 miles per hour) observed blowing from the south on October 17, 1940, over Rapid City, S. Dak., at an altitude of 1,594 meters (about 1 mile). A velocity of 62.8 m. p. s. (140 miles per hour) is the highest observed in the last 5 years in the layer from 2,500 meters to 5,000 meters. This wind was observed on October 24, 1939, blowing from the northwest at an altitude of 4,240 meters (about 2.6 miles) over Hartford, Conn. During this same 5-year period a still higher wind velocity 78.0 m. p. s. (174 miles per hour) was observed in the layer above 5,000 meters. This wind was blowing from the southwest at an elevation of 7,960 meters (about 14.9 miles) over Denver, Colo., on October 17, 1938

Table 1.—Mean free-air barometric pressure in millibars, temperature in degrees centigrade, and relative humidities in percent, obtained by airplanes and radiosondes during October 1941

									iii pu				00mu	es uu		70.00	<i>Jei</i> 1	041										
										Statio	ons wit	h ele	vatio	ns in m	ieters a	bove	sea le	evel										
	Albı	iquero (1	ue, N. ,620 m.)	Mex.			- ita, Ga. 90 m.)		Bi		s, N. D 5 m.)	ak.		Boise (86	, Idaho 4 m.)		Bi		ville, To m.)	9 x.	:	Buffal (22	o, N. Y 1 m.)	7.	9	Charles (14	ton, 8. m.)	c.
Altitude (meters) m. s. l.	Number of ob-	Pressure	Temperature	Relative bu-	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob-	Pressure	Tenperature	Relative hu-
Surface	30 30 30 30 30 30 30 30 29 29 29 27 26 24 23 20 19 10	628 553	12. 6 9. 5 8. 8 -1. 1 166. 8 -12. 6 -19. 3 -26. 4 -33. 6 -47. 0 -52. 5 -66. 8 -64. 1 -66. 2	5 544 5 555 8 49 8 49 8 39 8 37 8 37	31 31 31 31 31 31 31 31 31 31 31	986 963 909 857 808 761 717 635 561 493 433 379 330 287 248 213 182 155 132 194 79 67 57	18. 0 15. 7 13. 5 11. 0 8. 6 3. 9 -1. 8 -20. 8 -28. 0 -35. 5 -43. 2 -50. 8 -57. 8 -68. 7	63 67 64 56 56 56 34 31 28 29 29 29 28	31 31 31 31 31 31 31 31 31 31	957 901 848 797 749 620 546 478 417 362 313 270 232 199 170 145 124 106 90 77 666	6. 0 8. 2 5. 7 3. 2 1. 4 -1. 2 -6. 0 -12. 4 -19. 2 -26. 5 -33. 8 -41. 0 -47. 1 -54. 1 -54. 1 -54. 8 -56. 2 -58. 0 -58. 0 -56. 9	65 63 61 57 55 51 50 49 48 47	28	917 902 850 850 752 707 623 547 418 363 314 270 232 199 170 145 105 90 76	2.6 -0.7 -7.1 -13.4 -20.3	66 62 61 61 61 59 55 54 53	31 31 31 31 31 31 31 31 30 29	1, 014 958 905 854 806 759 7156 634 561 494 434 380 288 248 214 183 155 131 110 93 78	24. 7 23. 4 20. 6 18. 4 16. 9 5. 0 -0. 7 -13. 9 -20. 7 -28. 1 -35. 8 -44. 0 -51. 9 -76. 6 -76. 6 -75. 8 -71. 6	71 68 59 56 55 53 46 43 41 40	31 31 31 31 31 31 30 30 30 30 28 26	992 960 903 850 752 707 623 549 481 367 318 275 237 203 147 116 107 91 77 77 66	10.5 8.4 6.4 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2	7 54 69 63 59 7 54 63 63 63 63 63 63 63 63 63 63	31 31 31 31 31 31 31 30 30 30 30 29 29 29	963 909 858 761 716 634 560 493 433 379 286 246 211 180 109 977	20.1 17.1 14.1 13.1 10.0 8.4.0 -15.1 -22.2 -29.2 -37.2 -45.4 -67.2 -75.1 -76.1 -76.1	00 77 733 77 76 77 88 66 88 44 88 44 88 22 22 23 24 24 24 24 24 24 24 24 24 24 24 24 24
					1 .						tations		elev			ers ab					TT		on. W.	37-	1	T-11-	4 TI	
Altitude		(1,61	r, Colo. 6 m.)				, Mich m.)	•	ļ	(1,19	o, Tex. 3 m.)			Ely, (1,90	8 m.)		. 1		lls, Mo 8 m.)				2 m.)			(178	t, Ill.	1.
(meters) m. s. l.	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Tenperature	Relative hu-
Surface	30 30 30 30 30 30 29 28 27 26 24 21 18 19 18 17 16 14	839 801 753 708 625 550 483 422 367 318 275 236 202 173 148 125 107 917	2. 4 -3. 4 -9. 6 -16. 3 -23. 2 -37. 5 -44. 6 -50. 4 -56. 8 -59. 2 -61. 2 -62. 4	64 63 64 63 59 58 52 49	31 31 31 31 31 31 30	994 958 903 850 800 752 707 625 550 423 369 3277 238 205 175 149 127 108 92 78	10. 4 11. 6 8. 6 7. 1 5. 0 2. 9 -1. 9 -20. 8 -14. 0 -27. 7 -35. 0 -41. 9 -53. 9 -63. 4 -63. 4	75 67 61 55 52 49 48 43 42 41 39	31 31 31 31 32 29 28 27 27 27 27 27 27 26 26 24 21 18	882 851 802 756 711 630 556 489 428 374 325 282 243 209 178 152 109 92 78 66	16. 6	55 57 56 50 41 40 38 38 37	31 31 31 31 31 31 31 31 31 30 30 30 30 30 29 29 26 15	809 752 707 624 548 480 420 364 272 234 200 171 146 91	3.7 5.6 4.1 .7 -5.4 -11.8 -18.7 -25.5 -30.0 -40.2 -46.1 -50.3 -57.0 -58.8 -60.4 -61.1	60 59 58 56 52 50 49	31 31 31 31 31 31 31 31 31 31 31 30 30 30 27 26 21	888 798 750 621 546 478 417 362 313 270 232 198 169 145 105 90	7. 2 7. 7 4. 8 1. 8 -1. 0 -6. 6 -12. 9 -19. 5 -26. 8 -41. 5 -47. 6 -52. 2 -57. 4 -58. 1 -58. 1	55 54 52 51 50 47 45 45	31 31 31 31 31 31 30 30 30 30 30 30 29 29 26 19 14	999 961 906 854 804 757 712 629 555 487 427 373 324 280 242 207 175 1128 108 92 77	13. 8 15. 3 13. 4 11. 2 9. 5 7 . 5 5 5 12. 3 - 12.	72 71 66 57 55 52 49 44 42 41 40 41	28 27 27 27 27	997 959 904 851 800 753 *708 025 551 484 423 369 320 277 238 204 174 148	11. 4 12. 1 9. 8 8. 0 6. 7 4. 9 2. 9 -2. 1 -21. 0 -27. 7 -34. 8 -41. 8 -48. 3 -54. 7 -60. 4 -64. 3	78 75 72 65 61 57 53 48 46 44 42 43

MONTHLY WEATHER REVIEW

Table 1.— Mean free-air barometric pressure in millibars, temperature in degrees centigrade, and relative humidities in percent, obtained by airplanes and radiosondes during October 1941—Continued

														ing Ot	tober	-0.7.			nava									
										8	tations	with	elev	ations	in mete	rs ab	ove s	ea leve	el									
	La	ke Ch (5	arles, I m.)	. 8.	Le	kebur (39	st, N. J m.)	1,1	Ŋ		d, Oreg m.)	•		Miam (4)	i, Fla. n.)		Na	shville (180	, Tenn m.)	•]	Norfoll (10	r, Va.¹ m.)	1		Oaklan (2	d, Cali m.)	if.
Altitude (members) m. s. l.	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu-
8urface	30 30 30 30 30 30 30 30 29 29 29 29 28 28 28 28 27 25 21 19 13 5	1, 016 960 906 855 806 760 716 634 560 493 433 287 248 213 183 155 132 111 93 78 66	-64. 6 -69. 6	78 74 69 65 56 51 48 42 40 39 38	32 32 32 32 32 32 32 32 31 30 29 27 26 4 22 22 22 19 18 18 16 10 5	1, 014 960 905 862 802 754 709 626 552 484 424 278 239 205 176 150 128 109 93 79 68	12. 0 12. 4 10. 2 8. 8 8. 8 77. 4 -13. 6 -20. 2 -27. 0 -34. 1 -47. 6 -53. 8 -62. 4 -63. 5 -62. 4 -63. 5	56 53 57 61 62 60	31 31 31 31 31 31 31 31	970 958 903 850 800 752 707 624 549 481 420 365 316 273 234 200 171 146 124 108 90	11. 2 12. 2 11. 4 8. 5. 7 3. 1 0. 4 -4. 7 -10. 5 -24. 8 -32. 1 -39. 3 -45. 9 -51. 1 -64. 9 -57. 7 -60. 1 -60. 6 -60. 6	66 69 69 64 59 44 43 43 42 39	31 31 31 31 31 31 30 30 30 30 29 29 29 29 29 28 21 4	1, 017 961 907 856 807 760 716 634 434 434 434 1380 331 249 214 183 156 132 111 94 78 66	24. 4 22. 5 19. 2 16. 2 13. 8 11. 9 4. 5 -0. 6 -13. 1 -20. 2 -27. 7 -35. 2 -42. 6 -50. 6 -58. 3 -71. 7 -75. 5 -75. 6	79 76 64 56 45 38 33 30 29	31 31	998 962 907 855 806 759 715 558 491 376 327 284 245 211 180 153 130 110 93 78	18. 4 19. 1 16. 6 14. 6 12. 4 2. 1 -3. 2 -9. 5 -9. 5 -3. 2 3. 2 3. 4 5. 3 -52. 8 -67. 3 -68. 5 -68. 7 -64. 1	66 69 66 62 59 54 46 44 42 39 36 33	22 22 22 22 22 22 22 21 21 18	1, 022 964 909 857 760 715 632 558	18. 2 18. 3 16. 6 13. 5 11. 5 9, 7 7. 1 1. 8 —5. 8	68 64 62 57 52 46 38	31 31 31 31 31 31 31 31 31 29 29 28 25 5 17 17 15 9 5	1, 015 957 902 850 753 708 625 551 483 422 367 318 276 237 204 174 149 127 108	6.1 3.4 -2.8 -15.8 -22.8 -29.6 -36.3 -42.3 -51.8 -55.1 -57.6 -60.8	4 58 7 51 7 49 4 46 1 40 33 33 7 30 8 32 8 32 8 32 8 32 8 32 8 32 8 32 8 32
													_							_	_	_						
										8	tations	with	elev	ations	in mete	ers al	bove s	sea lev	el						1		······	
A 3424 3 -	0	klaho Okla.	ma Cit; (391 m.)	y,)	(a, Nebr	•	F	ensaco	stations ola, Fla m.)				x, Ariz.		1	ortland	el l, Mair m.)	18	6	St. Lou (171	iis, Mo . m.)) .	8	t. Pau (22)	ıl, Min 5 m.)	n.
Altitude (meters) m. s. l.	Number of ob-	Pressure	ma Cit; (391 m.))					-d _s	ensaco (24	ola, Fla m.)	,ı	-qo	Phoeni	x, Ariz.		r of ob-	ortland	i, Mair		Number of ob- servations	Pressure	rature E	e hu-	r of ob-	Pressure	Tenperature (.w. fr	Relative hu-

See footnotes at end of table.

Table 1.—Mean free-air barometric pressure in millibars, temperature in degrees centigrade, and relative humidities in percent, obtained by airplanes and radiosondes during October 1941—Continued

						rpian	ies ar	ia rac	uoson —	aes	auri	ng O	ctoo	er 19	41	Cont	inued	ا 								
									Stati	одз W	rith e	levati	ons ir	met	ers abo	- 076 868	level									
	Sar	1 Anto	nio, Te m.)	x.)	Sa	n Dieg (19:	o, Cal m.)	if.1	Sault	Ste. (22	Mari 1 m.)	e, Mic	eh.	s	eattle (27	, Wash m.)	.1	6	Spoka (5	ne, Wa 98 m.)	sh.		Wash	ingto (25 m	n, D. 1.)	c.
Altitude (meters) m. s. l.	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of 6b- servations	Pressure	Temperatura	Relative hu-	midity	Number of ob- servations	Pressure	Temperature	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob-	Progenite	omeou i	Temperature	Relative hu- midity
Surface	31 31 31 31 31 31 31 31	996 960 960 855 806 759 716 634 560 494 434 331 288 249 214 184 157 133 112 95 80 68	21. 1 19. 1 17. 0 14. 4 12. 0 9. 8 4. 8 -0. 8 -13. 1 -13. 1 -19. 8 -26. 4 -33. 8 -41. 2 -48. 7 -55. 9 -62. 2 -67. 6 -71. 1 -71. 1 -69. 1	90 85 79 79 80 67 58 53 48 45 41	300 300 300 300 299 289 286 277 24 233 211 19 16 12 10 7	1, 011 956 901 849 800 753 708 626 552 486 552 425 371 322 278 240 206 177 151 128	-26. 5 -33. 0 -39. 7 -45. 5 -51. 3 -56. 0 -59. 7	63 54 46 42 36 40 37 39	311 311 311 311 311 312 26 26 25 24 23 223 221 20 18 155 9	999 955 90 844 7997 744 7707 614 366 311; 277 141 122 110; 88 76	70 44 76 11 68 02 -77 -12 68 02 -78 -12 6 -18 6 -28 6 -28 -32 -32 -52 -52 -52 -52 -53 -58 -58 -58 -58 -58 -58 -58 -58	3. 7 5. 7 5. 7 7. 1 7. 1 7. 1 7. 1 7. 1 7. 1 7. 1 7		311 311 311 311 311 311 311 311 299 299 277 277 227 221 19 14	1, 015 959 903 850 799 751 705 621 5457 416 361 312 269 231 198 169 144 1124 88 74	7.7 5.0 2.6 0.1	90 888 85 79 74 69 52 55 54 49	31 31 31 31 31 31 31 31 30 30 28 27 27 27 25 24 20 13 8	90 84 79 75 70 62 54 41 36 31 28 23 19 16 14 12 10 8	22 8. 99 5. 88 3. 0 0. 55 -2. -14. 7 -20. 6 -28. -36. 2 -43. 8 -49. 0 -53. 7 -55. 8 -56. 4 -57. 2 -58. 4 -57.	1 77 9 65 5 62 6 61 1 44 4 4 4 4 4 4 5 66 7 7	332223333333222222222222222222222222222	9 4 9 8 8 8 8 2 7 2 1 6 1 5 1 2	963 908 908 805 758 713 330 556 -128 -128 -178 -178 -109 -178 -1		
			-	A ncho	rage, A	laska	Atla	ntic Sta	ation N								bove se	ТТ		o Solo	C. Z.1	.	Fairt	nanks	s, Alas	
		_		(42 m.)			(3 1	n.)			(3	m.)		_	(7	m.)			(15 n	a.) ———	_	r an c	(156 1	m.)	
Altitude (mete	ers) m.	s. 1.	Number of ob-	Servations Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob- servations	Pressure	Temperature	Relative hu-	Number of ob-	Pressure	Тетрегатиге	Relative hu- midity	Number of ob- servations	Pressure	Temperature	Number of ob-	Servations	rressure	Temperature	Relative hu-
Surface 500 1,000				29 5 29 4 28 3 27 2 27 2 27 2 27 2 25 11 23 10 20 11 5 10	866 — (31 — 3886 — 1910 — 1925 — 225 — 225 — 226 — 3742 — 4494 — 4494 — 4886 — 4886 — 4886 — 48888 — 4888 — 488888 — 48888 — 48888 — 48888 — 48888 — 48888 — 48888 — 48888 — 48888 — 48888 — 48888 —	7. 4 63 3. 9 60 5. 8 59 7. 6 59 1. 2 1. 5 1. 5 1. 5 1. 5 1. 6 1. 6 1. 6 1. 7 1. 8 1. 8 1. 8 1. 9 1. 9	28 28 28 28 28 27	1, 017 960 905 852 802 755 710 628 426 426 427 323 280 242 207 151 128 109 92 78 66 66 66	-62. 1 -64 2	80 80 72 63 55 40 38 35 35 34	27 27 27 27 27 27 27 27 28 25 25 24 23 22 20 17 17 15 19 6	1, 017 960 904 852 802 754 710 627 552 425 370 205 175 149 126 107 91 77 65 55	-27. -34. -42. -49. -55. -60.	8 8 7 6 6 6 4 4 3 5 6 4 4 3 5 6 4 4 3 5 6 6 2 2 5 5 8 9 4	27 26 26 23 23 19	1, 003 944 887 832 7832 686 686 601 525 457 396 342 294 2217 186 159 118 108 75	1. 6 0. 2 -2. 3 -6. 8 -9. 6 -12. 5 -18. 3 -30. 9 -37. 9 -44. 0 -49. 1 -50. 8 -47. 8 -47. 8 -48. 0 -48. 4 -48. 9	79 72 66 66 62 63 62 58 56 55	19 19 19 19 19 19 6		26. 4 24. 9 22. 2 19. 4 16. 7 13. 5 10. 7 4. 1	92 79 78 70 66 64 60 58	30 30 30 30 30 30 30 30 30 29 29 28 26 26 26	731 684 598 522 453 392 290 290 213 183 156 134 115	-2.8 -4.6.4 -8.5 -10.4 -15.6 -21.9 -28.4 -34.9 -41.3 -51.3 -50.3 -50.4 -51.3 -50.4	75 77 78 76 74 72 70 67 67

See footnotes at end of table.

Table 1.—Mean free-air barometric pressure in millibars, temperature in degrees centigrade, and relative humidities in percent, obtained by airplanes and radiosondes during October 1941—Continued

		ir piu	ites u	iu ru	110801	iues u	iarinę	7 000	06/ 16	941	Com	muec								
							Stat	ion's w	ith ele	vations	s in me	eters ab	ove sea	a level						
	June	au, Al	aska (4	9 m.)	K	tchika (26	n, Ala m.)	ska	Non	ae, Ala	ska (1	4 m.)	St.	Thom:	as, V. n.)	I.1 2	San J	uan, F	P. R. (1	5 m.)
Altitude (meters) m. s. l.	Number of observations	Pressure	Temperature	Relative hu- midity	Number of observations	Pressure	Temperature	Relative humidity	Number of observations	Pressure	Temperature	Relative hu- midity	Number of observations	Pressure	Temperature	Relative hu-	Number of observations	Pressure	Temperature	Relative hu- midity
·· · · · · · · · · · · · · · · · · · ·	ngo Sqo	Pre	Ten	Rel	nN sqo	Pre	Ter	Bel	Na	P. P.	Ter	Rel	N See	Pre	Ter	Rel	ng o		Ter	Rel
Surface 500 1,000 1,000 1,500 2,000 2,000 3,000 4,100 5,000 6,000 7,000 8,000 10,000 11,000	5	462 401 346 298 256 220 189 162 139 120 102 88	1. 4 -1. 1 -3. 9 -6. 8 -9. 7 -15. 7 -22. 4 -28. 8 -35. 6 -42. 2 -47. 6 -48. 4 -47. 8 -47. 9 -48. 0 -48. 3 -48. 9	83 83 83 81 79 76 65 63	23 21 20 19 18 15 15 14 14 13 10 7	351 303 260 224 192 164 140 120	-7.5 -13.7 -19.9 -26.7 -33.3	88 88 88 86 80 76 72 70 68 66	30 30 29 29 29 29 29 28 26 26 24 22 20 20 18 17 17 17	886 831 7730 683 597 521 453 392 250 214 184 158 136 117 100 86	-4.5 -7.0 -9.2 -12.3 -15.0 -20.9 -26.8 -33.4 -45.5 -49.6 -50.6 -49.4 -47.1 -46.7 -46.5 -46.5	81 80 76 73 71 70 66 59 56 55	31 31 31 31 31 31		20. 8 17. 9 15. 5 13. 4 6. 0	93 78 75 71 60 49 36	31 31 31 31 30 30 29 28 28 28 28 28 28 28 28 28 28 28 28 28	248 214 184 156 133 112 94 80 67 57 48	17.3 14.6 12.3 9.9 4.5 -0.9 -6.8 -13.3 -20.0	

Late Reports for September and August

					ЭСР		er an		-8-											_
						£	Stations	with	elev	ations	in mete	rs at	ove s	ea leve	ol		_			
	8	ept em	BER 194	11							Αī	JGUS	г 194	1						
Altitude (meters) m. s. l.	St.	Thom:	as, V. I. m.)	. 1–2	3	Barrow (6	, Alask m.)	a		Joile (17	et, III. 8 m.)		St. I	homa: (8	s, V. I. m.)	1-2	87		and, W	. I.
	Number of observations	Pressure	Temperature	Relative hu- midity	Number of observations	Pressure	Temperature	Relative hu-	Number of observations	Pressure	Temperature	Relative humidity	Number of observations	Pressure	Temperature	Relative hu- midity	Number of observations	Pressure	Temperature	Relative hu- midity
Surface	26 26 26	1, 014 958 905 854 805 759 716 635	27. 8 24. 3 21. 6 18. 7 16. 5 14. 2 11. 8 6. 4	74 92 76 71 63 53 48 40	31 31 31 31 31 31 31 31 31 31 31 31 32 29 28 27 25 24 22 21 13 8	1, 016 957 900 846 795 747 701 617 542 474 413 358 309 266 6228 196 164 124 106 91 78 658	3.5 5.1 4.6 3.0 1.0 -1.2 -3.9 -9.5 -15.6 -22,2 -29.1 -36.6 -51.9 -44.0 -51.9 -48.0 -46.9 -46.2 -46.2	92 83 73 76 68 65 63 61 60 57 55 53 53	29 29 29 29 29 29 29 29 26 25 24 24 24 21 19 17 17 17 16 15	995 958 904 853 804 757 712 630 556 489 429 374 326 282 244 204 179 152 129 110 93 79	18. 5 20. 9 18. 4 15. 2 12. 3 9. 6 6. 9 1. 0 -4. 4 -10. 1 -16. 5 -23. 8 -31. 2 -38. 7 -46. 0 -52. 9 -58. 4 -62. 5 -64. 7 -63. 1 -59. 9	85 69 67 72 72 71 67 63 56 49 45 42 38	28 28 28 28 28 28 28 28 28 	1, 016 961 907 856 808 762 718 637	27. 8 24. 3 21. 3 18. 6 16. 4 14. 1 11. 8 6. 3	74 91 79 72 68 59 50 47	31 31 31 31 31 31 31 31 30 30 30 30 30 30 30 30 30 30 30 30 30	1, 013 958 905 854 806 759 716 634 560 494 434 331 288 214 183 155 131 111 93 78 66 56	27. 1 24. 3 21. 6 18. 8 16. 2 13. 6 10. 9 4. 8 -1. 0 -6. 9 -13. 3 -19. 9 -34. 5 -42. 7 -50. 9 -72. 7 -75. 7 -75. 7 -72. 6 -69. 1 -69. 5	

¹ U. S. Navy.

2 Airplane observations.

4 Observations made on Coast Guard vessels in or near the 5° square.
Lat. 35.00′ N. to 40.00′ N.
Long. 55.00′ W. to 60.00′ W.

4 Observations made on Coast Guard vessels in or near the 5° square.
Lat. 35.00′ N. to 40.00′ N.
Long. 45.00′ W. to 50.00′ W.

Note.—All observations taken at 11 p. m. 75th meridian time, except at Lakehurst, N.J., where they are taken near 5 a. m., E. S. T., at Norfolk, Va., where they are taken at about 6 a. m., and at Pearl Harbor, T. H., St. Thomas, V. I., and Coco Solo, C. Z., at about 7 a. m.

None of the means included in this table are based on less than 15 surface or 5 standard level observations.

Number of observations refers to pressure only, as temperature and humidity data are missing for some observations at certain levels; also, the humidity data are not used in daily observations when the temperature is below -40° C.

Table 2.—Free-air resultant winds based on pilot balloon observations made near 5 p. m. (75th meridian time) during October 1941. Directions given in degrees from North ($N=360^{\circ}$, $E=90^{\circ}$, $S=180^{\circ}$, $W=270^{\circ}$)—Velocities in meters per second

							-					-						, .							1		1100	1		=	1			1					_
	A (5	biler Tex 37 n	ne, n.)	que	buqi N. I 630 1	Mex.	Ì	Itani Ga. 199 m		1	illing Mont 095 n		N	mar . Da 512 m	k.		Boise Idah 366 m	ó	vil	row lle, 7 (7 m	ex.]	uffal N. Y 220 m			rling Vt. 132 n		Ch (arles S. C 17 m	ton, .)	1	hica Ili. 192 n		l	ocini Ohie 152 n	0 ′	ļ	enve Colo 627 1	١. `
Altitude (meters) m. s. l.	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity
Surface	29 28 26 26 22 20 14 13 11	253	5.0	30 30 24 23 19	198 195 207 228 256 254 256 267	3. 7 4. 0 5. 1 9. 7 13. 3 14. 0 20. 7	30 28	41 89 82 172 297 304 294 287 286 281 279 279	0.8	29 29 29 28 28 24 22 21 14	228 251 269 272 286 284 281 275 310	1.8 3.3 4.2 5.5 6.8 7.7 9.3 7.6	29 28 24 22 21 20 20 19	290 295 283 288 281 282 281 282 278 276	2. 2 2. 5 4. 7 6. 7 8. 2 10. 5 13. 3 16. 2 19. 3 21. 5	30 30 30 28 25 24 20 19	321 316 315 304 284 276 266 298 295 300	3. 2 3. 8 2. 6 2. 5 3. 2 2. 6 1. 8 2. 6 4. 7 8. 9	31 28 26 25 23 19 18 17 10	158 174	4. 1 4. 1 4. 5 4. 1	21 19 17 12 10	264 259 261 267 269 275 276 282	3. 5 5. 6 7. 8 9. 7 10. 8 13. 4 14. 9 15. 4	29 29 24 20 17 13 10	226 246 272 282 290 289 298	1. 4 4. 8 7. 4 9. 8 12. 0 13. 0 14. 5	31 31 29 27 27 27 28 25 22 12	129 125 131 212 252 266 277 276 272 273 284	1. 4 2. 2 1. 5 1. 5 2. 3 2. 8 4. 2 6. 6 7. 5 13. 6	29 27 24 23 23 22 18 17 14	273 276 284 289 295	1. 2 4. 0 6. 4 8. 4 10. 7 11. 7 17. 0 19. 5 25. 3 31. 0	26 25 24 22 21 17 16 11	243 243 248 255 269 267 268 284 285 276	0. 9 3. 4 5. 2 7. 6 8. 9 11. 3 12. 7 15. 3 18. 4 20. 9	29 28 24 22 21	8 42 260 270 266 267 263 268	0.9
	E (1,	Pa Tex 196	so, m.)		y, N ,910 i		Ju	3ran netic Colo ,413 r	on,	1	ensb N. C 71 m		1	Havr Mon 167 n	t.	vi.	ackso lle, I 14 m	Pla.		s Ve Nev 570 n	-		tle F Ark. 79 m			edfo Oreg 110 n	. 1	N (iam Fla. 10 m	ıi, .)	lis	innes Mi 265 n	nn.		Aobi Ala (8 m		1	shvi Fenr 94 m	1.
Altitude (meters) m. s. l.	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity
Surface	30 29 30 30 29 25 21 19 12	211 217 224 229 242 250 249 253 250	2.9	31 31 31 25 19 17 13	269 283 207 234 268 251 267 274 279 274	1. 6 3. 0 6. 4 7. 0 10. 6 14, 0	30 30 28 24 20 17	263 262 204 213 201 235 261	0.8 1.1 1.8 4.0 4.9 5.2 5.9	31 31 31	253 248 245 258 277 278 283 282 284 285 284 295	0. 8 2. 0 3. 5 4. 7 6. 7 8. 6 8. 9 10. 9 12. 4 16. 9 17. 4 22. 7	30 30 29 28 26 21 19 15 11	258 257 260 265 267 274 273 272 293 289	7. 3 7. 4 8. 1 9. 6	0.0	91 97 115 159 182 290 284 298 301 296 295 305 294	3.6 6.2 9.1	30 30 30 29 25 23 19 17 13 11		1. 1 1. 5 1. 3 1. 8 2. 2 1. 4 5. 0 8. 1 11. 7 18. 1 20. 3 22. 9		165 178 207 216 217 236 240 260 256	1. 9 2. 9 3. 3 4. 1 5. 7 6. 8 7. 7 9. 9 10. 8	29 29 29 28 27 27 24 24 23 21 18 15 10	326 318 344 282 180 258 298 327 332 342 327 314 324	0. 9 1. 3 1. 2 0. 1 0. 9 0. 4 3. 3 4. 5 6. 4 7. 6 14. 4 15. 3	31 31 30 26 25 24 24 25 23 18 16 12	95 93 89 91 102 127 114 108 83 72 63 336 297	4. 0 5. 3 4. 3 3. 1 2. 5 2. 7 3. 1 2. 7 2. 3 1. 7 2. 5 3. 4 4. 6	28 24 22 17 16 16 16 12 10		0. 9 1. 8 2. 9 4. 9 9. 1 11. 1 13. 6 16. 3 19. 7 21. 4 27. 6	31 31 29 26 24 19 15 14 11	115 117 128 70 101 273 282 245 283	1. 9 1. 6 1. 1 0. 2 0. 8 1. 4 2. 5	31 31 31 31 27 24 20 16 13 10	213 218 216 226 243 258 278 275 289	2.8 4.5 5.3 5.2 6.1 7.2
	1 1	w Y N. Y 15 m	7.		akla: Cali (8 m	f.	Ok Cit	laho ty, O 102 n	ma, kla. 1.)	1	mah Nebi 306 m	•	l	boer Ariz 338 n		8	pid (). Da)82 m	k.	1	. Lo Mo 181 I		tor	an A nio, T 180 m	n- Fex. 1.)	1	n Di Cali: 15 n			ault i Marie Mich 230 m	e, 1.	l '	eatt Wasi 12 m	h.	'	poka Wasi 603 n	h.	Wa (shing D. C 24 m	ton,
Altitude (meters) m. s. i.	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity
Surface		275 287 279 288 290 294 293	3. 4 5. 1 6. 2 8. 1 10. 6 14. 0	31 30 28 26 25 25 22 21 20 12	258 325 335 323 323 326 332 329 315 308 270	2.9 1.5 1.8 2.1 2.2 2.7 4.6 7.6 9.4 10.9 11.7	25 25 22 20 20 19 17 16 13 10	198 183 189 218 222 239 241 246 245 253	2. 2 2. 7 4. 6 7. 0 8. 8 9. 5 10. 0 13. 1 14. 3 18. 2	29 28 27 23 19 15 14 14 14 12	191 183 203 237 261 281 284 275 269 272 274	1. 5 2. 4 3. 2 4. 6 5. 4 6. 1 8. 9 10. 5 13. 0 14. 7 19. 9	31 31 31 31 30 30 29 26 21 15	200	0. 9 1. 2 1. 1 1. 9 3. 7 5. 7 6. 2 8. 8 11. 8 13. 0 16. 5	1 11	356 346 310 287 283 286 277 274 276 284	1. 8 3. 8 5. 7 6. 6 8. 7 10. 2 11. 7	27 26 25 22 21 19 18 15	197 211 210 238 264 267 268 280 277 287	1. 0 2. 3 4. 3 6. 0 7. 3 9. 6 10. 5 12. 5 15. 2 16. 5	30 30 30 28 26 22 20 15 12 11	120 130 146 169 186 206 222 237 242 242	2. 2 3. 2 3. 7 4. 5 5. 4 5. 9 5. 7 7. 0 8. 4 10. 6	2 31 31 30 5 29 25 24 21 19 14	250 247 222 169 226 267 277 289 267	2. 9 3. 0 0. 9 0. 6 0. 7 2. 6 4. 0 7. 5 10. 1	17 13 10	292 288 289 275 273 280	3. 2 3. 9 7. 8 8. 8 10. 1 11. 2	30 30 27 20 18 17 14 12	252	3. 5 5. 0 4. 5 3. 4 4. 1 4. 5	28 28 25 24 22 19 12 11	210 212 224 235 245 250 274 279	1. 5 2. 2 3. 5 4. 8 5. 4 6. 3 3. 3 2. 2	31	291	2.6 4.4 5.6 7.5 10.4 11.6 13.1 15.4 17.7 20.6 28.6

Table 3.—Maximum free air wind velocities, (m. p. s.), for different sections of the United States based on pilot-balloon observations during October 1941

		Surface	to 2,50	0 me	ters (m. s. l.)	:	Between 2,	500 and	5,000	meters (m. s. l.)		1	Above 5,)00 n	neters (m. s. l.)
Section	Maximum ve- locity	Direc- tion	Altitude (m.) m. s. l.	Date	Station	Maximum ve-	Direc- tion	Altitude (m.) m. s. l.	Date	Station	Maximum ve-	Direc-	Altitude (m.) m. s. l.	Date	Station
Northeast 1 East-Central 2 Southeast 3 North-Central 4 Central 5 South-Central 6 Northwest 7 West-Central 8 South-West 9 South-West 9	42. 6 36. 2 26. 3 39. 4 36. 6 34. 0 36. 0 33. 4 29. 9	{WSW WSW SW WSW SW WSW NNW	1,740 2,420 1,650 2,100 2,500 1,970 1,900 430	5 8 27 27 7 5 17 3 2	Sacramento, Calif	55. 4 28. 2 63. 2 45. 0 38. 4 37. 6 51. 6	NW NW NW SSW SW WSW NNE SW	4, 400 3, 940 3, 170 4, 970 3, 370 3, 770 4, 650	23 10 10 7 4 26 10 2 13 2	Kylertown, Pa. Huntington, W. Va. Spartanburg, S. C. Muskegon, Mich. Wichita, Kans. Oklahoma City, Okla. Butte, Mont. Reno, Nev. Winslow, Ariz. Bakersfield, Calif.	70.0 55.2 74.5 73.0	WNW W WNW WSW SW WNW		17 28 20 10 9 31 2 25 8	Albany, N. Y. Greensboro, N. C. Miami, Fla. Minneapolis, Minn. Omaha, Nebr. San Antonio, Tex. Great Falls, Mont. Reno, Nev. Las Vegas, Nev.

WEATHER ON THE NORTH ATLANTIC OCEAN

By H. C. HUNTER

Atmospheric pressure.—The pressure during October 1941, averaged above normal over large portions of the North Atlantic, especially near the coasts of the South Atlantic States, Cuba, and the Bahamas. On the other hand it averaged below normal near the Maritime Provinces, Newfoundland, and Labrador.

The extremes of pressure noted in vessel reports at hand were 1,036.2 and 991.5 millibars (30.60 and 29.28 inches). The high reading was noted during the forenoon of the 30th, about 50 miles from Montauk Point, in a south-southwest direction. Table 1 shows that the pressure was slightly higher at Halifax, Nova Scotia, that day. The low mark was noted near southeastern Newfoundland during the early afternoon of the 20th. During other portions of the month both Belle Isle and Halifax recorded lower pressures.

In low latitudes readings slightly below 1,000 millibars (29.53 inches) were noted by two vessels near 29° N., 75° W., on the 10th, when within the area affected by a storm of tropical origin. However, this storm, when crossing the Bahama Islands several days earlier, as indicated in an article on page 303 of this Review, resulted in a pressure of 964.4 millibars (28.48 inches) on Cat Island.

Table 1.—Averages, departures, and extremes of atmospheric pressure (sea level) at selected stations for the North Atlantic Ocean and its shores, October 1941

Station	Average pressure	Depar- ture from normal	Highest	Date	Lowest	Date
Horta, Azores. Belle Isle, Newfoundland. Halifax, Nova Scotia. Nantucket. Hatteras. Turks Island ¹ Key West New Orleans.	Millibars 1, 020. 0 1, 007. 9 1, 014. 8 1, 018. 0 1, 020. 3 1, 016. 0 1, 016. 3 1, 017. 3	Millibars +0.4 -3.3 -2.5 +.4 +2.3 +1.8 +2.4 +.4	Millibars 1, 031 1, 029 1, 038 1, 036 1, 030 1, 019 1, 022 1, 024	13 30 30 30 29 18 14	Millibars 1, 009 986 989 1, 000 1, 009 1, 012 1, 011 1, 010	7 12 11 10 10 16 6 2

¹ For 27 days.

Indiana, Illinois, Iowa, Nebraska, Kansas, and Missouri.
 Mississippi, Arkansas, Louisiana, Oklahoma, Texas (except El Paso), and Western

** Montana, Idaho, Washington, and Oregon.

** Wyoming, Colorado, Utah, Northern Nevada, and Northern California.

** Southern California, Southern Nevada, Arizona, New Mexico, and extreme West

Cyclones and gales.—The vessel reports available indicate a comparatively quiet October. In middle latitudes two strong gales and several fresh gales occurred. The period centering on the 20th appears to have been the most eventful, and from the 8th to 11th likewise was somewhat unquiet.

The opening week and the period from 22d to 26th seem to have been remarkably undisturbed from the 30th parallel northward.

Tropical disturbances.—In another portion of this issue an account is presented of two disturbances of tropical origin. The earlier of these cyclones was noted to northward of the Virgin Islands on the 3d; it moved westward across the Bahamas and extreme southern Florida, turned northward over the Gulf and traversed parts of western Florida, southern Georgia and South Carolina, then moved southeastward over the Atlantic to near latitude 30°, thence continued mainly eastward till beyond the field of

This storm was of comparatively small diameter during much of its course, and was for awhile remarkably intense, especially over the Bahamas. A few lives are reported lost in the Bahamas and some in northwestern Florida. No vessel report that relates to this storm indicates a higher wind than a strong gale (force 9).

The later Low was of short path and little moment. Its track was approximately the middle portion of the track of the earlier storm, and the time was about 13 days later. The greatest wind force noted in vessel reports as a result of this disturbance was a moderate gale (force 7).

Line squalls.—About 150 miles to eastward of the extreme southeastern coast of Maryland line squalls were noted from a vessel which furnishes this report:

October 1, 9:30 p. m., 75th meridian time (equivalent to October 2, 2:30 a. m., Greenwich mean time), in latitude 38°04′ N., longitude 73°00′ W., barometer 30.16 inches (1021.3 millibars), having risen 0.03 inch during 1½ hours preceding, temperature of air 72°, of water 70°; three line squalls, moving southeastward, ranging from about 15 to 20 miles in length, passed at intervals of about 5 to 8 minutes. The clouds appeared in a very compact mass and in an almost straight line. During this time the wind shifted from southalmost straight line. During this time the wind shifted from southwest, Beaufort force 3, to northwest, same force; within the succeeding 3½ hours the wind turned partly back, becoming west, force 2. Other clouds at the time of the squalls were alto-cumulus, about two-tenths of sky being covered, apart from the line squall masses which covered a third of the overhead as they passed. Sea small southwest. Ship's course north-northeast, speed 14 knots.

Maine, Vermont, New Hampshire, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania and Northern Ohio.
 Delaware, Maryland, Virginia, West Virginia, Southern Ohio, Kentucky, Eastern Tennessee and North Carolina.
 South Carolina, Georgia, Florida, and Alabama.
 Michigan, Wisconsin, Minnesota, North Dakota, and South Dakota.

Note.—All data based on available observations, departures compiled from best available normals related to times of observation, except Hatteras, Key West, Nantucket, and New Orleans, which are 24-hour corrected means.